

Equivalence of the four versions of Tsallis' statistics

G. L. Ferri¹, S. Martínez², and A. Plastino²

¹ *Facultad de Ciencias Exactas, Universidad Nacional de La Pampa, Argentina*

² *La Plata Physics Institute (IFLP)*

National University La Plata and Argentine National Research Council (CONICET)

C.C. 727 , 1900 La Plata, Argentina

Abstract

In spite of its undeniable success, there are still open questions regarding Tsallis' non-extensive statistical formalism, whose founding stone was laid in 1988 in JSTAT. Some of them are concerned with the so-called *normalization* problem of just how to evaluate expectation values. The Jaynes' MaxEnt approach for deriving statistical mechanics is based on the adoption of (1) a specific entropic functional form S and (2) physically appropriate constraints. The literature on non-extensive thermostatics has considered, in its historical evolution, four possible choices for the evaluation of expectation values: (i) 1988 Tsallis-original (TO), (ii) Curado-Tsallis (CT), (iii) Tsallis-Mendes-Plastino (TMP), and (iv) the same as (iii), but using centered operators as constraints (OLM). The 1988 was promptly abandoned and replaced, mostly with versions ii) and iii). We will here (a) show that the 1988 is as good as any of the others, (b) demonstrate that the four cases can be easily derived from just *one* (any) of them, i.e., the probability distribution function in each of these four instances may be evaluated with a unique formula, and (c) numerically analyze some consequences that emerge from these four choices.

PACS: 05.30.-d, 05.30.Jp

KEYWORDS: Tsallis Statistics, Thermodynamics.

I. INTRODUCTION

James Bernoulli published in the *Ars Conjectandi* the first formal attempt to deal with probabilities in 1713 and Laplace further formalized things in his *Théorie analytique des Probabilités* of 1820. As we know, in the intervening years Probability Theory (PT) has grown into a rich, powerful, and extremely useful branch of Mathematics. Contemporary Physics heavily relies on PT for a large part of its basic structure, Statistical Mechanics [1, 2, 3], of course, being a most conspicuous example. One of PT basic definitions is that of the mean value of an observable $\mathcal{A}$ (a measurable quantity), which in its original form it is taken as a *lineal combination* of the different probabilities that $\mathcal{A}$ adopts for the different accessible states (we denote them by the index i). Let A stand for the linear operator or dynamical variable associated with $\mathcal{A}$. Then

$$\langle A \rangle = \sum_i p_i A_i. \quad (1)$$

It is well known that, in some specific cases, it becomes necessary to use “weighted” mean values, of the form

$$\langle A \rangle = \sum_i f(p_i) A_i, \quad (2)$$

with f an analytical function of the p_i . This happens, for instance, when there is a set of states characterized by a distribution with a recognizable maximum and a large tail that contains low but finite probabilities. One faces then the need of making a pragmatical (usually of experimental origin) decision regarding f .

Nonextensive thermostatistics (NET)

Tsallis’ nonextensive thermostatistics (NET) is nowadays considered by many authors to be a new paradigm for Statistical Mechanics (SM) [4, 5]. It encompasses a whole *family* of SM-formulations, all of them based upon Tsallis’ nonextensive information measure [4, 5]

$$S_q = k_B \frac{1 - \sum_{i=1}^W p_i^q}{q - 1}, \quad (3)$$

where k_B stands for Boltzmann’s constant and $\{p_n\}$ is a set of normalized probabilities. The real parameter q is called the index of nonextensivity. The literature on Tsallis’ thermo-

statistics considers three possible choices for the evaluation of mean- or expectation-values (EVs) within the nonextensive scenario. Since EVs are always regarded as constraints in the associated MaxEnt approach [6], three types of q -statistics will thereby ensue. Consider the physical quantity O that, in the microstate n ($n = 1, 2, \dots, W$), adopts the value o_n . Let p_n stands for the microscopic probability for such microstate. The O -EV is evaluated according to three distinct recipes, that we illustrate below with reference to Gibbs' canonical ensemble, where it is assumed that the a priori physical information is given by the mean value of a Hamiltonian ($O \equiv H$) of spectrum $\{o_i \equiv \epsilon_i\}$, and there are two Lagrange multipliers, one associated to normalization (α) and the other to the mean energy (inverse temperature β).

i) First choice: Tsallis (1988) [7]

$$\langle O \rangle = \sum_{i=1}^W p_i o_i. \quad (4)$$

The associated, Tsallis canonical distribution was (see the pertinent demonstration below with reference to Eq. (21))

$$p_i = [(1 - q)(\alpha + \beta \epsilon_i)/q]^{1/(q-1)}. \quad (5)$$

The awkward location of the multiplier α makes p_i hard to evaluate, save for very special cases like that of the harmonic oscillator [7]. This motivated people to look for better alternatives (see below). Other normalization choices thus emerged that typically employ “weighted” mean values and avoid the explicit presence of the multiplier α :

ii) Second choice: Curado - Tsallis (1991) [8]

$$\langle O \rangle = \sum_{i=1}^W p_i^q o_i, \quad (6)$$

with a canonical distribution given by

$$\begin{aligned} p_i &= (Z_q)^{-1} [1 - (1 - q) \beta \epsilon_i]^{1/(1-q)} \\ Z_q &= \sum_i [1 - (1 - q) \beta \epsilon_i]^{1/(1-q)}. \end{aligned} \quad (7)$$

iii) Third choice: Tsallis - Mendes - Plastino (1998) [9]

$$\langle O \rangle = \frac{\sum_{i=1}^W p_i^q o_i}{\sum_{i=1}^W p_i^q}. \quad (8)$$

We need to define here a normalization factor (NOT a Lagrange multiplier)

$$\mathcal{X}_q = \sum_i p_i^q. \quad (9)$$

The pertinent canonical distribution, that is *self-referential* because the p_i depends upon $\mathcal{X}_q$. Since it reads [9]

$$\begin{aligned} p_i &= Z_q^{-1} \left[1 - \frac{(1-q)\beta}{\mathcal{X}_q} (\varepsilon_i - U_q) \right]^{1/(1-q)} \\ \mathcal{X}_q &= (Z_q)^{1-q} \quad (\text{characteristic identity}) \\ Z_q &= \sum_i \left[1 - \frac{(1-q)\beta}{\mathcal{X}_q} (\varepsilon_i - U_q) \right]^{1/(1-q)}, \end{aligned} \quad (10)$$

this means that even if one knows U_q (as a real number) and the spectrum (we call this the canonical practitioner's input), this is not enough to obtain straightforwardly the probability distribution. One still needs to repeatedly iterate (10) until numerical self-consistency is achieved, which can be difficult in practice. In order to overcome such a problem still another choice can be made, the so-called OLM one [10, 11, 12], which avoids the self-reference problem. The corresponding probability distribution arises from solving anyone of two different variational problems [13], namely, those arising after

1. replacing S_q by Rényi's information measure S^R

$$S^R = \frac{\ln \sum_{i=1}^W p_i^q}{1-q}, \quad (11)$$

and keeping the TMP constraints. We call this procedure the TMP-R one.

2. Maintain Tsallis' entropy, but replace the TMP constraints by "centered" mean values.

We call this procedure the OLM-one.

In both cases one finds [14]

$$\begin{aligned} p_i &= Z_q^{-1} [1 - (1-q)\beta (\varepsilon_i - U_q)]^{1/(1-q)} \\ \mathcal{X}_q &= Z_q^{(1-q)} \quad (\text{characteristic identity}) \\ Z_q &= \sum_i [1 - (1-q)\beta (\varepsilon_i - U_q)]^{1/(1-q)}, \end{aligned} \quad (12)$$

and it should be noted that, with access to the canonical practitioner's input, no numerical self-consistency treatment is now required.

Reasons for using weighted mean values

In the first stage of NET development, its pioneer practitioners made the pragmatic choice of using “weighted” mean values, of rather unfamiliar appearance for many physicists. Why? The reasons were of theoretical origin. It was at the time believed that, using the familiar linear, unbiased mean values, one was unable to get rid of the Lagrange multiplier associated to probability-normalization (see below), usually denoted with the letter α in the Literature. Since the Tsallis formalism yields, in the limit $q \rightarrow 1$, the orthodox Jaynes-Shannon treatment, the natural choice was to construct weighted EVs using the index q

$$\langle A \rangle_q = \sum_i p_i^q A_i, \quad (13)$$

the so-called Curado-Tsallis unbiased mean values (MV) [8]. As shown by Plastino and Plastino [15], employing the Curado-Tsallis (CT) mean values allowed one to obtain an *analytical* expression for the partition function out of the concomitant MaxEnt process [15]. This EV choice leads to a non extensive formalism endowed with interesting features: i) the above mentioned property of its partition function Z , ii) a numerical treatment that is relatively simple, and iii) proper results in the limit $q \rightarrow 1$. It has, unfortunately, the drawback of exhibiting un-normalized mean values, i.e., $\langle \langle 1 \rangle \rangle_q \neq 1$. The latter problem was circumvented in the subsequent work of Tsallis-Mendes-Plastino (TMP) [9], that “normalized” the CT treatment by employing mean values of the form

$$\langle A \rangle_q = \sum_i \frac{p_i^q}{\mathcal{X}_q} A_i. \quad (14)$$

Today, this is the orthodox normalization procedure for NET practitioners. However, the concomitant treatment is not at all simple. Numerical complications often ensue, which has encouraged the development of different, alternative lines of NET research. See, for instance (a by no means exhaustive list), [16, 17, 18, 19, 20, 21, 22] and references therein.

Present goals

In this work we revisit the MaxEnt approach with unbiased mean values as prior knowledge, and show that *it is possible* to obtain a probability distribution that i) does not explicitly

contain the obnoxious α factor, ii) allows a seemingly analytical expression for the partition function out of the concomitant MaxEnt process (although it remains self-referential), iii) it is normalized, and iv) leads to the proper result in the $q \rightarrow 1$ limit. It does exhibit some numerical drawbacks, similar to those faced by the TMP treatment, but without the theoretical limitations that affect both the CT and the TMP formalisms. Since the treatment here advanced allows one to avoid weighted expectation values, it might be preferable to both the CT and and TMP ones on Ockham's razor-grounds.

Moreover, we will show that all four choices described above, namely, (4), (6), (8), and OLM, are *equivalent* in the sense that the four cases can be easily derived from just *one* (any) of them, i.e., the probability distribution function in each of these four instances may be evaluated with a *unique* formula, that (in the canonical instance) reads, for a given q -value

$$p_i = \frac{[1 - (1 - q^*) \beta^* \varepsilon_i]^{1/(1-q^*)}}{\sum_i [1 - (1 - q^*) \beta^* \varepsilon_i]^{1/(1-q^*)}} \\ \equiv \frac{\exp_q(-\beta^* \varepsilon_i)}{\sum_i \exp_q(-\beta^* \varepsilon_i)}; \text{ for } \beta^* \varepsilon_i < \frac{1}{1 - q^*}, \quad (15a)$$

$$p_i = 0; \text{ if } \beta^* \varepsilon_i \geq \frac{1}{1 - q^*}, \quad (15b)$$

where β^* is an appropriate pseudo-inverse temperature ($= 1/k_B T^*$) which depends on β , $\exp_q$ is the well-known q -generalized exponential [4]

$$\exp_q(x) \equiv [1 + (1 - q)x]^{1/(1-q)}, \quad (16)$$

and q^* is an effective nonextensivity index, suitably related to q (see Table 1). The extension of (15) to the general case of M observables $\mathcal{O}_j$, ($j = 1, \dots, M$), that involves M Lagrange multipliers λ_j , ($j = 1, \dots, M$), is straightforward.

II. RESCUING TSALLIS 1988 VERSION

Classical canonical treatment

For didactic purposes we will restrict ourselves first to the canonical ensemble and the classical formulation. The forthcoming section will treat things in quantal fashion for general

ensembles. The Boltzmann constant will be taken as unity in this Section. We write the internal energy (the mean value of the Hamiltonian) as the linear probabilities combination

$$U_q = \langle H \rangle = \sum_{i=1}^W p_i H_i, \quad (17)$$

where the sum is taken over the W micro-states of the system. The MaxEnt procedure entails maximizing Tsallis information measure subject to the constraints given by the internal energy and normalization of the probabilities

$$\sum_i p_i = 1. \quad (18)$$

The quantity to be maximized is

$$L_T = S_q - \alpha \sum_i p_i - \beta U_q, \quad (19)$$

i.e.,

$$\frac{\partial L_T}{\partial p_i} = \frac{q}{1-q} p_i^{q-1} - \alpha - \beta H_i = 0, \quad (20)$$

wherefrom we easily find

$$p_i = \left[\frac{(1-q)}{q} (\alpha + \beta H_i) \right]^{1/(q-1)}. \quad (21)$$

This is the result obtained and used in the very first paper on the Tsallis formalism [7], and subsequently *abandoned* i) because of the presence of the α term in (21) and ii) due to the fact that it was seemingly impossible to obtain a proper partition function. The step that transforms the above treatment into a useful one is performed via a simple trick. Note that we can obtain α by first multiplying Eq. (20) by p_i and then summing over i , which yields

$$\alpha = \frac{q}{1-q} \mathcal{X}_q - \beta U_q. \quad (22)$$

Once we have α , the problem is solved. If we place α into Eq. (21) we obtain

$$p_i = \left[\mathcal{X}_q - \frac{q-1}{q} \beta H_i \right]^{1/(q-1)}, \quad (23)$$

where $\delta H_i = H_i - U_q$. To explicitly obtain the partition function is now a simple task. Using Eq. (18) one immediately finds

$$1 = \frac{\sum_i \left[1 - \frac{q-1}{q} \frac{\beta}{\mathcal{X}_q} \delta H_i \right]^{1/(q-1)}}{\mathcal{X}_q^{1/(1-q)}}, \quad (24)$$

so that, if we agree to call

$$\bar{Z}_q = \sum_i \left[1 - \frac{q-1}{q} \frac{\beta}{\mathcal{X}_q} \delta H_i \right]^{1/(q-1)}, \quad (25)$$

we obtain the well-known, typical TMP result [9]

$$\mathcal{X}_q = \bar{Z}_q^{1-q}, \quad (26)$$

and

$$p_i = \frac{\left[1 - (q-1) \frac{\beta}{q\mathcal{X}_q} \delta H_i \right]^{1/(q-1)}}{\bar{Z}_q}. \quad (27)$$

It is clear that the need of having a real probability forces one to retain only the positive values of the expression between brackets (cut-off condition). Notice that the result (26) resembles the TMP one and that the presence of the quantity $\mathcal{X}_q$ makes the probability distribution a self referential one, although the proper limit when $q \rightarrow 1$ is indeed obtained

$$p_i|_{q=1} = \frac{e^{\beta U} e^{-\beta H_i}}{e^{\beta U} \text{Tre}^{-\beta H_i}} = \frac{e^{-\beta H_i}}{\text{Tre}^{-\beta H_i}}. \quad (28)$$

It remains as yet to ascertain whether the entropy is indeed a maximum, a point that has not been sufficiently discussed thus far. Taking the second derivative of L_T we easily obtain

$$\frac{\partial^2 L_T}{\partial p_i^2} = -q p_i^{q-2}, \quad (29)$$

i.e., for $q > 0$ the entropy will be a maximum. *This result entails that negative q -values are not acceptable.*

General quantal treatment of the Tsallis' 1988 version

Within the general quantum nonextensive framework, all information concerning the system of interest is provided by the statistical operator $\hat{\rho}$. To find it one has to extremize the

information measure

$$S_q = \frac{\mathcal{X}_q - 1}{1 - q}, \quad (30)$$

where now

$$\mathcal{X}_q = \text{Tr} (\hat{\rho}^q), \quad (31)$$

subject to the normalization requirement

$$\text{Tr} (\hat{\rho}) = 1 \quad (32)$$

and the assumed a priori knowledge of the expectation values (EV)s of, say, M relevant observables $\hat{O}_j$ with $j = 1 \dots M$

$$\langle \hat{O}_j \rangle = \text{Tr} (\hat{\rho} \hat{O}_j). \quad (33)$$

The density or statistical operator $\hat{\rho}$ is here the quantum version of the set of probabilities of the preceding Subsection. Following exactly the same procedure as before, with M Lagrange multipliers λ_j , we obtain a density operator of the form

$$\hat{\rho} = \frac{\hat{f}^{1/(q-1)}}{\bar{Z}_q}, \quad (34)$$

with a partition function given by

$$\bar{Z}_q = \text{Tr} \hat{f}^{1/(q-1)}, \quad (35)$$

where

$$\hat{f} = 1 - (q - 1) \sum_j \frac{\lambda_j}{q \mathcal{X}_q} \delta \hat{O}_j \quad (36)$$

is sometimes called the configurational characteristics. It is finite only if its argument is positive. $\hat{f}$ is null if its argument is negative (cut-off condition). The quantities

$$\delta \hat{O}_j = \hat{O}_j - \langle \hat{O}_j \rangle_q \quad (37)$$

are usually called centered operators.

Connections between de 1988 and the TMP formalism

Comparing Eq. (34) with the density operator of TMP treatment [9], we see that it is easy to pass from one to other via an appropriate mapping. If the subindex u stands for the 1988 NET version we have

$$\begin{aligned}
q - 1|_u &\mapsto 1 - q|_{TMP} \\
\lambda_j/q|_u &\mapsto \lambda_j|_{TMP}.
\end{aligned}
\tag{38}$$

An interesting consequence of Eq. (38) ensues: as S_q will be convex for $q|_u > 0$ we discover that the convex region for S_q within the TMP formalism will be $q|_{TMP} < 2$. Now, it has been shown in Ref. [10, 11] that in the TMP instance one also finds the requirement $q > 0$. Thus, TMP and 1988 results should be valid for $q \in (0, 2)$.

III. EQUIVALENCE OF THE DIFFERENT TSALLIS VERSIONS

As stated above one takes, for a numerical treatment of the canonical ensemble i) the internal energy U_q as a numerical datum and ii) the energy spectrum. This provides one with a probability distribution. We have called i) and ii) the canonical practitioner's input. We advance here an *alternative route* (we call it the *parametric approach*), in which, *instead* of giving U_q as a numerical input data, one a priori fixes the numerical value of an effective inverse temperature β^* . With such an alternative procedure one is able to achieve two different goals, *for a given energy-spectrum*:

- a unification of all Tsallis' techniques, from a theoretical viewpoint, and
- a very convenient numerical procedure, from a practical perspective.

Indeed, by recourse to an analysis similar to the one given above, effected now for *all 4 types* of Tsallis distribution mentioned in the Introduction, it is easy to show that this parametric technique can i) link the four NET versions and ii) give the pertinent four distribution-forms a “universal” (common) appearance. The ensuing results are summarized in Table 1 for the canonical ensemble. The energy spectrum $\{\varepsilon_i\}$ constitutes the basic a priori data. A power-law distribution-form (common to all 4 Tsallis versions) of exponent $q^*(q)$ can be built up, for any q -value, that reads

$$p_i = Z_q^{-1} \exp_{q^*(q)}(-\beta^* \varepsilon_i); \text{ whose (version - dependent) ingredients are :} \tag{39}$$

- an effective inverse temperature β^* , given a priori as a number,

- Z_q , the normalization constant (partition function), and
- an “effective” nonextensivity index $q^* = q^*(q)$, given in the Table. Indeed, save for the case of the 1988 version, we have $q^* = q$.

In all four cases, U_q is evaluated a posteriori, as indicated in the Table. One can demonstrate that the appropriate entropic form S_q is (a posteriori) maximized by (39) (for *this* particular U_q). Also,

1. T is the “true” thermodynamic temperature that a thermometer would measure, and
2. β is the Lagrange multiplier of the putative MaxEnt process one could have employed to reach the “parametric” PD (39) using the canonical practitioner’s input.

By TO we mean Tsallis’ 1988 original treatment, and by TMP-R the OLM one [10, 11, 12]. *From the canonical practitioner’s viewpoint* β^* is, save for the CT and OLM (TMP-R) instances, a rather formidable quantity that makes the probability distribution a self-referential one (the signature of the self-referential character is the appearance of $\mathcal{X}_q$ in the expression for β^*). *This is not so from the parametric perspective.* As explained below, if one fixes β^* at the outset, considering it as an input parameter, the self-referential difficulty vanishes. One is then able to numerically build in fast fashion the PD (39) for very many β^* -values in the interval $[0, \infty]$ and then “read” off the Table the appropriate values for $T(\beta)$, U_q , $\mathcal{X}_q$ associated to any given β^* .

Table 1: Non-extensive Statistics				
Probability Distribution Function				
$p_i = Z^{-1} \exp_{q^*}(-\beta^* \varepsilon_i)$, if $\beta^* \varepsilon_i < \frac{1}{1-q^*}$; $Z = \sum_i \exp_{q^*}(-\beta^* \varepsilon_i)$				
$p_i = 0$, if $\beta^* \varepsilon_i > \frac{1}{1-q^*}$				
	β^*	q^*	U_q	T
<i>TO</i>	$\beta^* = \frac{\beta}{q\chi_q + (q-1)\beta U_q^{TO}}$	$q^* = 2 - q$	$U_q^{TO} = \sum_i p_i \varepsilon_i$	$k_B T = \frac{k_B T^* - (q-1)U_q^{TO}}{q\chi_q}$
<i>CT</i>	$\beta^* = \beta$	$q^* = q$	$U_q^{CT} = \sum_i p_i^q \varepsilon_i$	$k_B T = k_B T^*$
<i>TMP</i>	$\beta^* = \frac{\beta}{\chi_q + (1-q)\beta U_q^{TMP}}$	$q^* = q$	$U_q^{TMP} = \frac{\sum_i p_i^q \varepsilon_i}{\sum_i p_i^q}$	$k_B T = \frac{k_B T^* - (1-q)U_q^{TMP}}{\chi_q}$
<i>TMP-R</i>	$\beta^* = \frac{\beta}{1 + (1-q)\beta U_q^{TMP-R}}$	$q^* = q$	$U_q^{TMP-R} = \frac{\sum_i p_i^q \varepsilon_i}{\sum_i p_i^q}$	$k_B T = k_B T^* - (1-q)U_q^{TMP-R}$

Notice that, if any one of the four Tsallis' probability distributions (PDs) is available, we can with it, using the appropriate equations of the version at hand, evaluate the quantities U_q , $\mathcal{X}_q$, and from them obtain T . Now for translating into another Tsallis' version we

- use our PD to evaluate the U_q -value of the new version. With it and
- the above values for $\mathcal{X}_q$ and T , compute the *new* β^* , appropriate for this second version.
- Finally, employ Eq. (39).

In order to gain some insight onto the properties of each Tsallis-version we have computed, for two different q values, namely, $q = 0.6$ (Figs. 1 and 3) and $q = 1.5$ (Figs. 2 and 4) the relative internal energy $U_q/(h\nu)$ (Figs. 1 and 2) and the specific heat C_q/k_B (Figs. 3 and 4) for a system of equally spaced levels labelled by a "quantum number" i , like in the case of the harmonic oscillator, for instance (inter-level distance $= \varepsilon_0$), with $i = 0, 1, 2, \dots$ and $\varepsilon_0 = h\nu$. For all four Figs., the horizontal axis represents $k_B T/(h\nu)$ and the $q = 1$ results are also shown for comparison. A glance to Figs. 1-4 shows that the four Tsallis versions yield different results, and that the OLM instance is the one that resembles the orthodox $q = 1$ situation in closer fashion. In particular, notice in Fig. 2 that the OLM- $q = 1.5$ curve is identical to that for $q = 1$. Specific heat undulations are a consequence of Tsallis' cut-off, as explained in Ref. [12]. The *asymptotic* behavior when $T \rightarrow \infty$ for U_q and C_q is given by the following expressions:

$$U_q \sim T^\alpha$$

$$C_q \sim T^{\alpha-1}$$

where the appropriate α -exponent values, for each Tsallis version, are shown below:

TO	CT	TMP	$TMP - R$
$\alpha = \frac{1}{q}$	$\alpha = 2 - q$	$\alpha = \frac{1}{q}$	$\alpha = 1$

Advantages of the parametric form (39)

It should be remarked that using the probability distribution (PD) (39) exhibits the advantage of avoiding self-reference no matter *which* of the normalization choices one uses. In writing down the PD in the fashion

$$p_i = Z_q^{-1} \exp_{q^*}(-\beta^* \varepsilon_i); \quad Z_q = \sum_i \exp_{q^*}(-\beta^* \varepsilon_i), \quad (40)$$

you can numerically input, for a given

$$\begin{aligned} q^* &= 2 - q; \text{ for the 1988 version} \\ q^* &= q; \text{ for all the other versions,} \end{aligned} \tag{41}$$

any β^* -value whatsoever and you get the PD. No self-reference here. Since the data are the spectrum, β^* , and q^* , we speak of a “parametric” calculation. With such a PD you evaluate both $\mathcal{X}$ and U_q and then proceed to ascertain the value of the physical temperature T using the relations given in Table 1. From a numerical viewpoint therefore, (40) provides one with a convenient tool to work out nonextensive thermostatistics problems.

CONCLUSIONS

We have revisited the four different extant versions of Tsallis’ nonextensive thermostatistics and shown that they can be unified (Cf. Table 1) in the sense of deriving the transformations that provide a link between any given pair of versions. It must be emphasized, however, that *numerical results are different* for the distinct versions. In particular, we have shown that the currently abandoned original Tsallis version of 1988 is a quite legitimate one.

-
- [1] R.K. Pathria, *Statistical Mechanics* (Pergamon Press, Exeter, 1993).
 - [2] F. Reif, *Statistical and thermal physics* (McGraw-Hill, NY, 1965).
 - [3] J. J.Sakurai, *Modern quantum mechanics* (Benjamin, Menlo Park, Ca., 1985).
 - [4] M. Gell-Mann and C. Tsallis, Eds. *Nonextensive Entropy: Interdisciplinary applications* (Oxford University Press, Oxford, 2004); V. Latora, A. Rapisarda, and C. Tsallis, *Physica A* **305** (2002) 129 (and references therein); S. Abe and Y. Okamoto, Eds. *Nonextensive statistical mechanics and its applications* (Springer Verlag, Berlin, 2001); V.Latora, A.Rapisarda, and C. Tsallis, “Fingerprints of nonextensive thermodynamics in a long-range Hamiltonian system”, 2002 *Physica A* **305** 129 (and references therein); S. Abe and Y. Okamoto, Eds. *Nonextensive statistical mechanics and its applications* Springer Verlag, Berlin, 2001; A. R. Plastino and A. Plastino, “From Gibbs microcanonical ensemble to Tsallis generalized canonical distribution” 1994 *Phys. Lett. A*, **193** 140. A periodically updated bibliography on nonextensive thermostatistics can be found in the URL “<http://tsallis.cat.cbpf.br/biblio.htm>”.

- [5] C. Tsallis, “Nonextensive Statistics: Theoretical, Experimental and Computational Evidences and Connections”, 1999 *Braz. J. of Phys.*, **29** 1; A. Plastino and A. R. Plastino, “Tsallis Entropy and Jaynes’ Information Theory Formalism”, 1999 *Braz. J. of Phys.*, **29** 50.
- [6] E. T. Jaynes, *Phys. Rev.* **106** (1957) 620; **108** (1957) 171; *Papers on probability, statistics and statistical physics*, edited by R. D. Rosenkrantz (Reidel, Dordrecht, Boston, 1987); A. Katz, *Principles of Statistical Mechanics, The Information Theory Approach* (Freeman and Co., San Francisco, 1967).
- [7] C. Tsallis, “Possible generalization of Boltzmann-Gibbs statistics”, 1988 *J. Stat. Phys.* **52** 479.
- [8] E. M. F. Curado and C. Tsallis, “Generalized statistical mechanics: connection with thermodynamics”, 1991 *J. Phys. A*, **24** L69.
- [9] C. Tsallis, R. S. Mendes, and A. R. Plastino, “The role of constraints within generalized nonextensive statistics”, 1998 *Physica A* **261** 534.
- [10] S. Martínez, F. Nicolás, F. Pennini and A. Plastino, “Tsallis’ entropy maximization procedure revisited”, 2000 *Physica A* **286** 489.
- [11] E. K. Lenzi, R. S. Mendes and L. R. da Silva, “Statistical mechanics based on Renyi entropy”, 2000 *Physica A* **280** 337.
- [12] G. Ferri, S. Martinez, and A. Plastino, “The role of constraints in Tsallis’ nonextensive treatment revisited”, 2005 *Physica A* **347** 205.
- [13] S. Martinez and A. Plastino, “Nonextensive treatment of Clausius equation”, 2005 *Physica A* **345** 493.
- [14] As a matter of fact, it can be shown that the basic relation $dU_q/dT = TdS_q/dT$ is valid in the TMP-R case but is violated by the OLM-treatment, notwithstanding the fact that in both instances the *same* probability distribution ensues, as shown in [13].
- [15] A. R. Plastino and A. Plastino, “Tsallis’ entropy, Ehrenfest theorem and information theory”, 1993 *Phys. Lett. A* **177** 177.
- [16] Sumiyoshi Abe, S. Martínez, F. Pennini and A. Plastino, “Nonextensive thermodynamic relations”, 2001 *Phys. Lett. A* **281** 126.
- [17] S. Abe, S. Martinez, F. Pennini and A. Plastino, “Classical gas in nonextensive optimal Lagrange multipliers formalism”, 2001 *Phys. Lett. A* **278** 249.
- [18] M. P. Almeida, “Generalized entropies from first principles”, 2001 *Physica A* **300** 424.

- [19] E. Vives and A. Planes, “Is Tsallis Thermodynamics Nonextensive?”, 2002 *Phys. Rev. Lett.* **88** 020601.
- [20] R.P. Di Sisto, S. Martínez, R.B. Orellana, A.R. Plastino and A. Plastino, “General thermostastical formalisms, invariance under uniform spectrum translations, and Tsallis q-additivity”, 1999 *Physica A* **265** 590.
- [21] R. Toral, “On the definition of physical temperature and pressure for nonextensive thermostatics”, 2003 *Physica A* **317** 209.
- [22] T. Yamano, “On the laws of thermodynamics from the escort average and on the uniqueness of statistical factors”, 2003 *Phys. Lett. A* **308** 364.

FIGURE CAPTIONS

Fig. 1: Relative internal energy $U_q/h\nu$ vs. $k_B T/h\nu$ ($q = 0.6$) for the four different instances of Tsallis-normalization (see text).

Fig. 2: Relative internal energy $U_q/h\nu$ vs. $k_B T/h\nu$ ($q = 1.5$) for the four different instances of Tsallis-normalization (see text).

Fig. 3: Specific heat C_q/k_B vs. $k_B T/h\nu$ ($q = 0.6$) for the four different instances of Tsallis-normalization (see text).

Fig. 4: Specific heat C_q/k_B vs. $k_B T/h\nu$ ($q = 1.5$) for the four different instances of Tsallis-normalization (see text).







